

REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>						
1. REPORT DATE (DD-MM-YYYY) 12/01/2015		2. REPORT TYPE Final		3. DATES COVERED (From - To) Jul 2012 - Sep 2013		
4. TITLE AND SUBTITLE Determining the molecular and genetic basis for diabetes in Navy bottlenose dolphins (<i>Tursiops truncatus</i>)				5a. CONTRACT NUMBER NA		
				5b. GRANT NUMBER N000141210617		
				5c. PROGRAM ELEMENT NUMBER NA		
				5d. PROJECT NUMBER NA		
6. AUTHOR(S) Montminy, Marc R.				5e. TASK NUMBER NA		
				5f. WORK UNIT NUMBER NA		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Salk Institute for Biological Studies 10010 North Torrey Pines Road La Jolla, CA 92037-1002				8. PERFORMING ORGANIZATION REPORT NUMBER NA		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 875 North Randolph Street Arlington, VA 22203-1995				10. SPONSOR/MONITOR'S ACRONYM(S) ONR		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) NA		
12. DISTRIBUTION/AVAILABILITY STATEMENT Unlimited						
13. SUPPLEMENTARY NOTES NA						
14. ABSTRACT CREB-ZF functions as a transcriptional repressor, at least in terms of its effects on gluconeogenic genes. Efforts are underway to determine whether CREB-ZF directly interferes with activation of the cAMP-responsive factor CREB or the CRTC family of CREB coactivators. CREB-ZF has been reported to modulate the unfolded protein response (UPR) although the underlying mechanism is unclear. Because it contains a leucine zipper DNA binding domain, CREB-ZF would be expected to compete for binding to CREB binding sites. We will test whether CREB-ZF displaces CREB and CRTC2 from gluconeogenic promoters and thereby reduces hepatic glucose production.						
15. SUBJECT TERMS Gluconeogenesis, CREB ZF, Fasting, Diabetes						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT	b. ABSTRACT	c. THIS PAGE			Marc R. Montminy	
U	U	U	UU	11	19b. TELEPHONE NUMBER (Include area code) 858-453-4100 x1394	

20150414072

**ONR Technical / Final Report:
July 2012 - September 2013**

PI Name: Marc Montminy, M.D., Ph.D.

Organization: Salk Institute for Biological Studies

ONR Award Number: N000141210617

Award Title: Determining the molecular and genetic basis for diabetes in Navy
bottlenose dolphins (*Tursiops truncatus*)

Contents

I. Scientific and Technical Objectives	3
II. Approach.....	3
III. Concise Accomplishments	4
IV. Expanded Accomplishments	4
V. Future Plans.....	8
VI. Major Problems/Issues.....	8
VII. Technology Transfer.....	8
VIII. Foreign Collaborations and Supported Foreign Nationals.....	8
IX. Productivity	9
A. Refereed journal articles	9
B. Non-refereed significant publications	9
C. Books and chapters	9
D. Technical reports	9
E. Patents.....	9
F. Awards/honors.....	9
X. Award Participants	9
Literature Cited.....	10

I. Scientific and Technical Objectives

Navy bottlenose dolphins (*Tursiops truncatus*) are susceptible to insulin resistance, and they appear to have a switch that turns a diabetes-like state on and off.^{1,2} While this metabolic state was initially thought to be normal for dolphins, recent studies have discovered that dolphins are susceptible to a number of chronic conditions associated with insulin resistance, including fatty liver disease, iron overload, urate nephrolithiasis, hyperlipidemia and chronic inflammation.³⁻⁶ These chronic conditions are present among Navy dolphins, and the study's primary hypothesis is that there is a molecular and genetic basis for insulin resistance in Navy dolphins. The scientific objective of this study was to begin characterizing the molecular and genetic basis of the gluconeogenic program in dolphins, with a focus on the genes and promoters for glucose-6-phosphatase (G6Pase) and PEPCK, two rate-limiting enzymes for gluconeogenesis in mammals. The technical objectives of this 1-year study were as follows:

- Amplify PCR genomic fragments corresponding to promoters for dolphin G6Pase and PEPCK genes
- Construct G6Pase and PEPCK promoter luciferase reporter constructs
- Use RT-PCR to obtain cDNAs for dolphin CREB-ZF, a member of the basic leucine zipper domain (bZIP) family of transcription factors with homology to cAMP response element binding protein (CREB)
- Perform transient transfection assays to test effects of CREB ZF over-expression on gluconeogenic gene expression in response to fasting signals

II. Approach

- A. Amplify PCR genomic fragments corresponding to promoters for dolphin G6Pase and PEPCK genes. The publically available partial dolphin genome from Baylor College of Medicine, based upon DNA from a Navy dolphin, was used to identify genomic fragments corresponding to promoters for dolphin G6Pase and PEPCK genes. Polymerase chain reactions (PCR) were used to amplify these fragments. To assess how well they were conserved, these dolphin gene sequences were compared to those from humans, mice, and rats. Targeted binding sites, including TATA boxes, included those for the transcription factors, forkhead box O1 binding protein (FOXO1) and CREB.
- B. Construct G6Pase and PEPCK promoter luciferase reporter constructs. Luciferase was used to assess successful transcriptional activity of dolphin G6Pase and PEPCK promoters. Cellular levels of cAMP were raised with forskolin (FSK) to mimic the fasting state, and relative luciferase activity was measured at 0, 2, 4, and 6 hours.
- C. Obtain cDNAs for dolphin CREB-ZF. While searching for the dolphin CREB gene, dolphin CREB-ZF was identified as a potential transcription factor. CREB-ZF is a member of the basic leucine zipper domain (bZIP) family of transcription factors with homology to CREB. RT-PCR was used to obtain cDNAs for dolphin CREB-ZF.
- D. Test effects of CREB-ZF over-expression on gluconeogenic gene expression. Transient transfection assays were used to test the effects of CREB-ZP on gluconeogenic gene expression in response to fasting signals (cAMP).

III. Concise Accomplishments

- A. Amplify PCR genomic fragments corresponding to promoters for dolphin G6Pase and PEPCK genes
 - Basic mechanisms for activation of gluconeogenic genes are conserved in dolphins.
- B. Construct G6Pase and PEPCK promoter luciferase reporter constructs
 - Dolphin PEPCK transcription increased in the face of increasing cAMP, supporting that this enzyme induces gluconeogenesis during the fasting state.
- C. Obtain cDNAs for dolphin CREB-ZF
 - Dolphin CREB-ZF, closely related to CREB, was identified as a potential transcription factor (genetic switch).
- D. Test effects of CREB-ZF over-expression on gluconeogenic gene expression
 - Dolphin CREB-ZF is a novel, negative regulator of gluconeogenesis.

Over the last few months of this grant, we continued our efforts to address the mechanism by which the transcription factor CREB –ZF regulates hepatic gluconeogenesis. Pilot studies revealed that CREB-ZF functions as a transcriptional repressor, at least in terms of its effects on gluconeogenic genes. Efforts are underway to determine whether CREB-ZF directly interferes with activation of the cAMP-responsive factor CREB or the CRTC family of CREB coactivators.

CREB-ZF has been reported to modulate the unfolded protein response (UPR) although the underlying mechanism is unclear. Because it contains a leucine zipper DNA binding domain, CREB-ZF would be expected to compete for binding to CREB binding sites. We will test whether CREB-ZF displaces CREB and CRTC2 from gluconeogenic promoters and thereby reduces hepatic glucose production. In future work, we will obtain full-length clones for dolphin CREB-ZF and test the effect of CREB-ZF on CREB/CRTC2 occupancy and recruitment.

IV. Expanded Accomplishments

- A. Amplify PCR genomic fragments corresponding to promoters for dolphin G6Pase and PEPCK genes. Figures 1 and 2 demonstrate the homology of these dolphin genes with those in human, mouse, and rat. FOXO1 and CREB binding sites are conserved in dolphin G6Pase promoter compared to human, mouse, and rat genes. CREB binding site is conserved on dolphin PEPCK promoter compared to human, mouse, and rate genes.

Figure 1. FOXO and CREB binding sites are conserved in dolphin G6Pase promoter compared to human, mouse, and rat genes.

	Foxo Binding Site	Foxo Binding Site	CREB Binding Site	
human	CTGTTTTT	GTGTGCC	TGTTTTT	CTATTTCACGTAAATCACCCCTGAACATGTTTGCATCAA
dolphin	CTGTTTTT	CTGTGCC	TGTTTTT	CCTATTTCACGTAAATCACACTGAACACGTTTGCATCAA
mouse	CTGTTTTT	GTGTGCC	TGTTTTT	GCTATTTCACGTAAATCACCCCTGAACATGTTTGCATCAA
rat	CTGTTTTT	GTGTGCC	TGTTTTT	GCTATTTCACGTAAATCACCCCTGAACATGTTTGCATCAA
	*****	*****	*****	*****
human	CCTACTGGTGATGCACCTTTGATCAATACATTTTAGACAAACGTGGTTTTT-GAGTCCAA			
dolphin	CCTACTGGTGATGCACCTTTGATCAATAGATTTTAGACAAAAGCGGTTTTT-GAGTCCAA			
mouse	CCTACTGATGATGCACCTTTGATCAATAGATTTTAGACAAAAGTGGTTTTTTGAGTCCAA			
rat	CCTACTGATGATGCACCTTTGATCAATAGATTTTAGACAAAAGTGGTTTTTTGAGTCCAA			


	TATA Box			
human	AGATCAGGGCTGGGTTGACCTGAATACTGGATACAGGGCATATAAAACAGGGGCAAGGCA			
dolphin	AGATCAGGGCTGGGTTGACCTGCAGACTGGATACAGAGTGTATAAAACAGAGGCAAGACA			
mouse	AGATCAGGGCTGGATTGACCTACAGACTGAATCCAGGGCATATAAAACAGGGGCAAGGCA			
rat	AGATCAGGGCTAGGTTGACCTACAGACTGAATCCAGGGCATATAAAAT--GGGCAAGGCA			
	*****	*	*****	*****

CREB Binding Site

```
human      CAAAGCATAACTGACCCTGGCCGTGATCCAGAGACCTGCCCCCTGACGTCAGTGCGCAGC
dolphin    CAAAGCATAACTGACCGTGGCCGTGATCCAAAGACCAGCCCC-TGACGTCAGAGATGAGC
mouse      CAAACCGTG-CTGACCATGGCTATGATCCAAAGGCCTGCCCC-TTACGTCAGAGGCGAGC
rat        CAAACCGTG-CTGACCATGGCTATGATCCAAAGGCCGGCCCC-TTACGTCAGAGGCGAGC
          **** * *  *****  *****  *****  ** **  *****  *  *****  *  ****

human      CTCCTGGGTGCAGCTGAGGGGCAGGGC----TATTCTTTTCCACAGTATTTAAAGCTGG
dolphin    CTCCTTGGGTGTGGCTGAGGGGCTGGAC----CTCTGTCTTCATGGAATTTAAAGCCAG
mouse      CTCC--GGGTCCAGCTGAGGGGCAGGGCTGTCTCCTCCCTCTATATAGTATTTAAAGCAAG
rat        CTCC--AGGTCCAGCTGAGGGGCAGGGCTGTCTCCTCCCTCTGTATACATTTAAAGCGAG
          *****  ***  *****  * * *  *****  *  *  *****  *
```

Figure 3. Induction of PEPCK luciferase reporter in response to cAMP agonist (FSK)




C. Obtain cDNAs for dolphin CREB-ZF. Dolphin CREB-ZF had a basic leucine zipper (bZIP) domain, indicating that it is a transcription factor, or switch that may regulate gene activity. Dolphin CREB-ZF had distinct regions that were not homologous to CREB-ZF in mice, supporting that it may be a different construct.

Figure 4. Dolphin CREB-ZF sequence, including a bZIP domain, compared to mouse.

mouse	MRHSLTKLLAASGRDFPSRRDSREPPATRAPPREPSGAAAGAE-TPRPGSPDREQPHGDG
dolphin	MRHSLTKLLAASGSDSPTRSESPAPVATCSLPPDLTRAAAEDEGTAAAGSPGRKQPRGD- ***** * *: * * * : * : : * * * * * . * * . * * : * * : * *
mouse	DGGEPEARSGSRGSVAVRAPAPSPLKMEEEEDAIAMVPKEEPEDMDFLSGLELADLLDP
dolphin	-EGESDAGRGGRGIVAAAPSP-----EEMEEEAIASVPGEETEDMDFLSGLELADLLDP * * . : * * * . * * * * . * * : * * * * * * * * . * * * * * * * * * * * *
mouse	RQPDWHLEPGLSSPGPLSSSGGSESGLLRGDDDDTAAAEQRFSDLLQRLNIGIGGC
dolphin	RQPDWHLEPGLNSPGPLSSSGGSDSGGLWRGDDDDDEAAAAEQRFSDLLQRLNIGIGGC ***** . * * * * * * * * * : * * * * * * * * * : * * * * * * * * * * * *
mouse	SSGGDRGGGEKRRRKSPGAGGGG-ANDGNQAATKSPRKAARLNRLKKKEYVMGLES
dolphin	SSGSDSGGGEKRRRKSPGGGGSSGNDNNQAATKSPRKAARLNRLKKKEYVMGLES * * . * * * * * * * * * . * * . * * . * * * * * * * * * * * * * * * * * *
mouse	RVRGLAAENQELRAENRELGKRVQALQEEESRYLRAVLANETGLARLLSRLSGVGLRLTTS
dolphin	RVRGLAAENQELRAENRELGKRVQALQEEESRYLRAVLANETGLARLLSRLSGVGLRLTTS *****
mouse	LFRDSPAGDHDYALPVGKQPPEPREEDDAAGGVCLHVDKDKVSVEFCSACARKASSSLKM
dolphin	LFRDSPAGDHDYALPVGKQQQDLLEEDDSAGGVCLHVDKDKVSVEFCSACARKASSSLKM ***** : * * * : * * * * * * * * * * * * * * * * * *

D. Effects of CREB-ZF over-expression on gluconeogenic gene expression. Dolphin CREB-ZF over-expression inhibited PEPCK promoter activity in the face of elevated cAMP (Figure 5). This is the first report of CREB-ZF involvement in metabolism. Dolphin CREB-ZF is a novel, negative regulator and potential 'off switch' for gluconeogenesis.

Figure 5. Effect of CREBZF over-expression on dolphin pck1 promoter activity in cells exposed to cAMP agonist FSK for 4 hours. UT, untreated.



V. Future Plans (new white paper)

1. Obtain cDNA clones for dolphin CREB1 using RT-PCR.
2. Continue characterizing dolphin gluconeogenic program by evaluating relevant molecules and genes in liver, muscle, and adipose tissue.
3. Test the role of CREB-ZF in modulating hepatic gluconeogenesis by adenoviral over-expression or RNAi-mediated knockdown in mouse hepatocytes and in liver.
4. Determine the mechanism by which CREB-ZF inhibits gluconeogenic gene expression. Specifically, we will test whether CREB-ZF associates with CREB1 or with CREB-cofactors such as the CRTC.
5. Identify signals that regulate CREB-ZF expression and activity in liver. In particular, we will test effects of high fat diet feeding on CREB-ZF expression and activity. Does high fat diet feeding increase gluconeogenic gene expression by reducing CREB-ZF expression?
6. Test relative activities of mouse and dolphin CREB-ZF proteins. Could disparities in fasting glucose production between dolphins and other mammals reflect differences in respective CREB-ZF proteins?

VI. Major Problems/Issues

Lack of a full dolphin genome sequence led to difficulties in finding the transcription factor, CREB.

VII. Technology Transfer

Due to well conserved mechanisms for gluconeogenesis, drugs like metformin, which down-regulate gluconeogenic gene expression, may be effective in controlling fasting glucose in Navy dolphins. The data also suggest that increasing expression of the inhibitor CREB-ZF in hepatocytes may reduce gluconeogenic gene expression and circulating blood glucose levels in Navy dolphins.

VIII. Foreign Collaborations and Supported Foreign Nationals

None

IX. Productivity

A. Refereed journal articles

None

B. Non-refereed significant publications

None

C. Books and chapters

No books or chapters have been generated in relation to this project.

D. Technical reports

No technical reports have been generated in relation to this project.

E. Patents

No patents have been generated in relation to this project.

F. Awards/honors

No awards or honors have been generated in relation to this project.

X. Award Participants

The following non-military personnel received salary support from this ONR award during the reporting period: Bing Luan, Ph.D.

Literature Cited

1. Venn-Watson S, Ridgway SH (2007) Big brains and blood glucose: Common ground for diabetes mellitus in humans and healthy dolphins. *Comp Med* 57(4):390-5.
2. Venn-Watson S, Carlin K, Ridgway S (2011) Dolphins as animal models for type 2 diabetes: Sustained, postprandial hyperglycemia and hyperinsulinemia. *Gen Comp Endocrin* 170:193-199.
3. Venn-Watson S, Benham C, Carlin K, DeRienzo D, St. Leger J (2012) Hemochromatosis and fatty change: building evidence for insulin resistance in bottlenose dolphins (*Tursiops truncatus*). *J Zoo Wildl Med* 43(3 Suppl):S35-S47.
4. Venn-Watson S, Smith CR, Daniels R, Townsend F (2010) Clinical relevance of urate nephrolithiasis in bottlenose dolphins (*Tursiops truncatus*) *Dis Aqua Org* 89:167-177.
5. Venn-Watson S, Smith CR, Gomez F, Jensen ED (2011) Physiology of aging among healthy, older bottlenose dolphins (*Tursiops truncatus*): comparisons with aging humans. *J Comp Phys B* 181:667-680.
6. Venn-Watson S, Townsend FI, Daniels R, Sweeney J, McBain J, Klatsky L, Hicks C, Staggs L, Rowles T, Schwacke L, Wells RS, Smith CR (2010) Hypocitraturia in Atlantic bottlenose dolphins (*Tursiops truncatus*): Assessing a potential risk factor for urate nephrolithiasis. *Comp Med* 60:149-153.